arXiv:1210.5120v1 [hep-ph] 18 Oct 2012

# One-loop effects on top pair production in the littlest Higgs model with T-parity at the LHC

Bingfang Yang[1,2] and Ning Liu[1*]

[1] *College of Physics & Information Engineering, Henan Normal University, Xinxiang 453007, China*

[2] *Basic Teaching Department, Jiaozuo University, Jiaozuo 454000, China*

## Abstract

We calculate the one-loop corrections to top pair($t\bar{t}$) production at the Large Hadron Collider(LHC) for $\sqrt{s}$ = 8 TeV and 14 TeV in the context of the Littlest Higgs model with T-parity(LHT). We find that the relative corrections to $t\bar{t}$ cross section can maximally reach 4.5% for $\sqrt{s} = 8$ TeV in the favorable parameter space. We also study other observables in $t\bar{t}$ production, such as top charge asymmetry, spin correlation and polarization asymmetry. We note that the spin correlation is sensitive to the loop corrections from LHT model, and can be optimally reduced by $-14.5\%$ for $\sqrt{s} = 8$ TeV.

PACS numbers: 14.65.Ha,12.15.Lk,12.60.-i,13.85.Lg

* corresponding author: wlln@mail.ustc.edu.cn

## I. INTRODUCTION

Due to a heavy mass, top quark has been widely considered as a window of unveiling the new physics at TeV-scale. In particularly, the recent anomalous top quark forward-backward asymmetry observed at the Tevatron [1] may be a strong hint of new physics beyond the Standard Model(SM) [2], although many measurements from the Tevatron are consistent with the SM predictions. Besides, because of the small statistics, the study of top-quark properties is limited at the Tevatron. As a new generation of top quark factory, the LHC will copiously generate the top events via top pair productions and single top productions, which provides a good opportunity to precisely test the top quark properties and to search the new physics signals[3].

Notwithstanding the SM have been confirmed by various experiments from the LEP to LHC, it still has some drawbacks, such as the hierarchy problem. In order to solve this problem, the little Higgs model was proposed [4], which is done by making the Higgs as a pseudo-Goldstone boson. The littlest Higgs (LH) model[5] is an economical approach which implements the idea of the little Higgs, however, this model suffers strong constraints from electro-weak precision tests [6]and will reintroduce the fine-tuning problem in the Higgs potential[7]. A feasible way to overcome these difficulties is to impose a discrete symmetry called T-parity[8], which prevents the tree-level contribution from the heavy gauge bosons to the electro-weak observables. The T-parity also forbids the interactions that induce the triplet scalars to develop the VEV in the LH model. This resulting model is referred to as the littlest Higgs model with T-parity (LHT). One aspect of the LHT phenomenology in top-quark sector is that the top quark can have interactions with the LHT particles, such as heavy gauge bosons, mirror fermions, and heavy quarks $T^{\pm}$. These new interactions can contribute to the $t\bar{t}$ production at the loop level.

So far, the new particle beyond the SM has not been discovered at the LHC, so the scale of the new physics has to be high. In this case, study of new particle at the loop level is becoming increasingly important. Furthermore, since the produced top quarks decay before the hadronization, the spin information of top quarks will be inherited and manifested by its decay daughters. Therefore, spin polarization and spin correlation of top quark can be used to probe the mechanisms of top quarks productions and decays[9], and serve as a probe to new physics[10]. In this paper, we study the LHT corrections to

the process $pp \to t\bar{t}$ at the LHC with $\sqrt{s} = 8$ TeV and 14 TeV.

This paper is organized as follows. In Sec.II we give a brief review of the LHT model related to our work. In Sec.III we calculate the (un)polarized $t\bar{t}$ production and the correlation of the observables in the LHT model. Finally, we give our conclusions in Sec.IV.

## II. A BRIEF REVIEW OF THE LHT MODEL

The LHT is a non-linear $\sigma$ model based on the coset space $SU(5)/SO(5)$, with the global group $SU(5)$ being spontaneously broken into $SO(5)$ by a $5 \times 5$ symmetric tensor at the scale $f \sim \mathcal{O}(TeV)$, the gauged subgroup $[SU(2) \times U(1)]^2$ of $SU(5)$ is broken into the SM gauge group $SU(2)_L \times U(1)_Y$. From the $SU(5)/SO(5)$ breaking, there arise 14 Goldstone bosons which transform under the electroweak gauge group as follows:

$$\mathbf{1_0} \oplus \mathbf{3_0} \oplus \mathbf{2_{1/2}} \oplus \mathbf{3_{\pm 1}}. \tag{1}$$

where the subscripts indicate the hypercharges. We can denote the fields in these four representations as $\eta, \omega, H$ and $\phi$, respectively. After EWSB, $H$ can be decomposed as $H = (-i\pi^+\sqrt{2}, (v + h + i\pi^0)/2)^T$. Explicitly, they are described by the "pion" matrix $\Pi$, given by

$$\Pi = \begin{pmatrix} -\frac{\omega^0}{2} - \frac{\eta}{\sqrt{20}} & -\frac{\omega^+}{\sqrt{2}} & -i\frac{\pi^+}{\sqrt{2}} & -i\phi^{++} & -i\frac{\phi^+}{\sqrt{2}} \\ -\frac{\omega^-}{\sqrt{2}} & \frac{\omega^0}{2} - \frac{\eta}{\sqrt{20}} & \frac{v+h+i\pi^0}{2} & -i\frac{\phi^+}{\sqrt{2}} & \frac{-i\phi^0+\phi^P}{\sqrt{2}} \\ i\frac{\pi^-}{\sqrt{2}} & \frac{v+h-i\pi^0}{2} & \sqrt{4/5}\eta & -i\frac{\pi^+}{\sqrt{2}} & \frac{v+h+i\pi^0}{2} \\ i\phi^{--} & i\frac{\phi^-}{\sqrt{2}} & i\frac{\pi^-}{\sqrt{2}} & -\frac{\omega^0}{2} - \frac{\eta}{\sqrt{20}} & -\frac{\omega^-}{\sqrt{2}} \\ i\frac{\phi^-}{\sqrt{2}} & \frac{i\phi^0+\phi^P}{\sqrt{2}} & \frac{v+h-i\pi^0}{2} & -\frac{\omega^+}{\sqrt{2}} & \frac{\omega^0}{2} - \frac{\eta}{\sqrt{20}} \end{pmatrix} \tag{2}$$

Where $\omega^\pm, \omega^0, \eta$ are eaten respectively by 4 new heavy gauge bosons $W_H^\pm, Z_H, A_H$ whose masses up to $\mathcal{O}(v^2/f^2)$ are given by

$$M_{W_H} = M_{Z_H} = gf(1 - \frac{v^2}{8f^2}), M_{A_H} = \frac{g'f}{\sqrt{5}}(1 - \frac{5v^2}{8f^2}) \tag{3}$$

with $g$ and $g'$ being the SM $SU(2)$ and $U(1)$ gauge couplings, respectively. In the 't Hooft-Feynman gauge, the would-be Goldstone-Boson mass is the same as its corresponding gauge boson.

When T-parity is implemented in the fermion sector of the model we require the existence of mirror partners for each of the original fermions. For each SM quark, a copy of mirror quark with T-odd quantum number is added. We denote them by $u_H^i, d_H^i$, where i= 1, 2, 3 are the generation index, whose masses up to $\mathcal{O}(v^2/f^2)$ are given by

$$m_{d_H^i} = \sqrt{2}\kappa_i f, m_{u_H^i} = m_{d_H^i}(1 - \frac{v^2}{8f^2}) \tag{4}$$

where $\kappa_i$ are the diagonalized Yukawa couplings of the mirror quarks.

In order to cancel the one-loop quadratic divergent radiative corrections to Higgs mass parameter induced by top quark, an additional heavy T-even partner of the top quark $T^+$ is introduced. The implementation of T-parity then requires its own mirror quark $T^-$, which is T-odd under T-parity. Their masses up to $\mathcal{O}(v^2/f^2)$ are given by

$$m_{T^+} = \frac{f}{v}\frac{m_t}{\sqrt{x_L(1-x_L)}}[1 + \frac{v^2}{f^2}(\frac{1}{3} - x_L(1-x_L))] \tag{5}$$

$$m_{T^-} = \frac{f}{v}\frac{m_t}{\sqrt{x_L}}[1 + \frac{v^2}{f^2}(\frac{1}{3} - \frac{1}{2}x_L(1-x_L))] \tag{6}$$

where $x_L$ is the mixing parameter between the SM top-quark $t$ and the new top-quark $T^+$.

In the LHT model, the flavor structure is richer than the one of the SM due to the presence of the mirror fermions and their weak interactions with the ordinary fermions[11]. The mirror quark sector exists two CKM-like unitary mixing matrices as follows:

$$V_{Hu}, V_{Hd} \tag{7}$$

Note that $V_{Hu}$ and $V_{Hd}$ are related through the SM CKM matrix:

$$V_{Hu}^\dagger V_{Hd} = V_{CKM}. \tag{8}$$

These mirror mixing matrices are involved in the flavor changing interactions between the SM fermions and the mirror fermions which are mediated by the T-odd gauge bosons $(W_H^\pm, Z_H, A_H)$ or T-odd Goldstone bosons$(\omega^\pm, \omega^0, \eta)$. One cannot completely turn off the new mixing effects except with a universally degenerate mass spectrum for the T-odd mirror fermions. The mixing matrix $V_{Hd}$ can be conveniently parameterized, we follow Ref.[12] to parameterize $V_{Hd}$ with three angles $\theta_{12}^d, \theta_{23}^d, \theta_{13}^d$ and three phases $\delta_{12}^d, \delta_{23}^d, \delta_{13}^d$ as

follows

$$V_{Hd} = \begin{pmatrix} c_{12}^d c_{13}^d & s_{12}^d c_{13}^d e^{-i\delta_{12}^d} & s_{13}^d e^{-i\delta_{13}^d} \\ -s_{12}^d c_{23}^d e^{i\delta_{12}^d} - c_{12}^d s_{23}^d s_{13}^d e^{i(\delta_{13}^d - \delta_{23}^d)} & c_{12}^d c_{23}^d - s_{12}^d s_{23}^d s_{13}^d e^{i(\delta_{13}^d - \delta_{12}^d - \delta_{23}^d)} & s_{23}^d c_{13}^d e^{-i\delta_{23}^d} \\ s_{12}^d s_{23}^d e^{i(\delta_{12}^d + \delta_{23}^d)} - c_{12}^d c_{23}^d s_{13}^d e^{i\delta_{13}^d} & -c_{12}^d s_{23}^d e^{i\delta_{23}^d} - s_{12}^d c_{23}^d s_{13}^d e^{i(\delta_{13}^d - \delta_{12}^d)} & c_{23}^d c_{13}^d \end{pmatrix} \quad (9)$$

## III. NUMERICAL RESULTS AND DISCUSSIONS

In our calculation, we neglect the higher order couplings between the scalar triplet $\Phi$ and top quark and the high order $\mathcal{O}(v^2/f^2)$ terms in the masses of new particles. In the 't Hooft-Feynman gauge, we use the dimensional regularization scheme to regulate the ultraviolet divergences in the virtual corrections and adopt the on-shell renormalization scheme to remove them. The relevant Feynman diagrams for process $gg \to t\bar{t}$ and $q\bar{q} \to t\bar{t}$ in the LHT are depicted in Fig.1. The black dot and grey ellipse appearing in Figs.1 represent the renormalized vertexes $\hat{\Gamma}^{\mu}_{gt\bar{t}}$, $\hat{\Gamma}^{\mu}_{gq\bar{q}}$ and top quark self-energy at one-loop level respectively, whose diagrams are displayed in Fig.2 and Fig.3. We list the explicit expressions of these amplitudes in Appendix. We analytically and numerically checked that the divergences in the renormalized vertex and propagator have been canceled. We also find that there are no divergences in the box diagrams. We use the parton distribution function CTEQ10[13] with renormalization scale and factorization scale $\mu_R = \mu_F = m_t$. The SM parameters input in our numerical calculations are taken as[14]

$$\begin{aligned} &G_F = 1.16637 \times 10^{-5}\ \text{GeV}^{-2}, \quad \sin^2\theta_{\text{W}} = 0.231, \\ &\alpha_e = 1/128,\ M_{Z_L} = 91.2\ \text{GeV},\ \text{m}_{\text{t}} = 172.9\ \text{GeV}. \end{aligned} \quad (10)$$

Recently the ATLAS and CMS experiments at the LHC have independently discovered a Higgs-like resonance with mass about 125 GeV[15], so we take $m_h = 125GeV$ in our numerical calculations. The relevant LHT parameters are the scale $f$, the mixing parameter $x_L$, the Yukawa couplings $\kappa_i$ and the parameters in the matrices $V_{H_u}, V_{H_d}$. For the mirror fermion masses, we get $m_{u_H^i} = m_{d_H^i}$ at $\mathcal{O}(v/f)$ and assume that the masses of the first two generations are degeneracy.

$$m_{u_H^1} = m_{u_H^2} = m_{d_H^1} = m_{d_H^2} = M_{12}, m_{u_H^3} = m_{d_H^3} = M_3 \quad (11)$$




FIG. 1: Feynman diagrams of the one-loop correction to the process $pp \to t\bar{t}$ in the LHT model. The black dot and grey ellipse represent the renormalized vertexes $\hat{\Gamma}^{\mu}_{gt\bar{t}}$, $\hat{\Gamma}^{\mu}_{gq\bar{q}}$ and top quark self-energy respectively, whose diagrams are displayed in Fig.2 and Fig.3.

For the matrices $V_{H_u}$ and $V_{H_d}$, we follow Ref.[16] to choose the following scenario: $V_{H_u} = 1, V_{H_d} = V_{CKM}$. In this scenario, the contribution of the mirror quarks will come entirely from the third family ones and the additional heavy quarks $T^+, T^-$. In our calculations, we consider the constraints in Refs.[17] and scan the parameter regions: $f = 500 \sim 2000$GeV, $x_L = 0.1 \sim 0.9$, $\sqrt{2}\kappa_i = 0.6 \sim 3$. Since the new parity violating interactions between top quark and LHT particles can not only affect the $t\bar{t}$ production rate but also the spin polarization, we will discuss the LHT corrections to the (un)polarized top pair production by using the following observables and their correlations.

(i) For the unpolarized $t\bar{t}$ production, we calculate the relative corrections for total $t\bar{t}$ production cross section($\delta\sigma/\sigma$), charge asymmetry($A_C$)[22] at the LHC and the top




FIG. 2: The effective $gt\bar{t}, gq\bar{q}$ vertex diagrams at one-loop level in the LHT model.




FIG. 3: The effective fermion propagator diagrams at one-loop level in the LHT model.

quark forward-backward asymmetry($A_{FB}^t$)[18] at the Tevatron.

$$\delta\sigma/\sigma = \frac{\sigma_{tot} - \sigma_{SM}}{\sigma_{SM}}, \tag{12}$$

$$A_C = \frac{\sigma(\Delta|\eta_t| > 0) - \sigma(\Delta|\eta_t| < 0)}{\sigma(\Delta|\eta_t| > 0) + \sigma(\Delta|\eta_t| < 0)}, \tag{13}$$

$$A_{FB}^t = \frac{\sigma(\Delta y_t > 0) - \sigma(\Delta y_t < 0)}{\sigma(\Delta y_t > 0) + \sigma(\Delta y_t < 0)}. \tag{14}$$

where $\Delta y_t$ ($\Delta\eta_t$) is the (pseudo)rapidity difference of the top and anti-top quark in the laboratory frame. The total cross section $\sigma_{tot}$ contains the contributions from both the SM tree-level and the interference between SM and the LHT model.

(ii) For the polarized $t\bar{t}$ production, we calculate the spin correlation($\delta C$)[9], the polarization asymmetry($P_t$)[19] and the left-right asymmetry($A_{LR}$)[19].

$$C = \frac{(\sigma_{RR} + \sigma_{LL}) - (\sigma_{RL} + \sigma_{LR})}{\sigma_{RR} + \sigma_{LL} + \sigma_{RL} + \sigma_{LR}}, \tag{15}$$

$$\delta C = \frac{C_{tot} - C_{SM}}{C_{SM}}, \tag{16}$$

$$P_t = \frac{(\sigma_{RL} + \sigma_{RR}) - (\sigma_{LR} + \sigma_{LL})}{\sigma_{RL} + \sigma_{RR} + \sigma_{LL} + \sigma_{LR}}, \tag{17}$$

$$A_{LR} = \frac{\sigma_{RL} - \sigma_{LR}}{\sigma_{RL} + \sigma_{LR}}. \tag{18}$$

Here, the subindices $L(R)$ represent left($\lambda_{t(\bar{t})} = -1/2$) and right-handed($\lambda_{t(\bar{t})} = +1/2$) top(antitop) quarks, respectively.

### A. Unpolarized top quark pair production




FIG. 4: The relative correction of the top-quark pair production cross section $\delta\sigma/\sigma$ as the function of $f, x_L$ for $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV, respectively.

In Fig.4, we show the relative correction $\delta\sigma/\sigma$ versus $f$ and $x_L$ at the LHC for $\sqrt{s} = 8, 14$ TeV respectively. We can see that the maximal value of the relative correction to the $t\bar{t}$ cross section can reach 4.5% for $\sqrt{s} = 8$ TeV and 1.6% for $\sqrt{s} = 14$ TeV, respectively. Because the LHC is a proton-proton machine, about 90% of the top pair production rate will come from the $gg$ initial state and the remaining 10% from $q\bar{q}$. Moreover, we find the relative correction mainly comes from the process $q\bar{q} \to t\bar{t}$, which agrees with the results in Ref.[20]. With the center-of-mass energy $\sqrt{s}$ increasing, the

production rate comes from the $gg$ increases faster so that the relative correction $\delta\sigma/\sigma$ for $\sqrt{s} = 14$ TeV is smaller than the one for $\sqrt{s} = 8$ TeV. With the scale $f$ increasing, the relative correction display the random behaviour, which is mainly because of the Yukawa couplings such as $\bar{u}^i_H \omega^0 u^j : \frac{g}{2M_{Z_H}}[m_{u^i_H} P_L - m_{u^j} P_R](V_{Hu})_{ij}$. Due to $M_{Z_H}$ and $m_{u^i_H}$ are both proportional to the scale $f$, so their effects will cancel each other out with the $f$ changing. The relative correction $\delta\sigma/\sigma$ decrease with $x_L$ increasing, which indicates that the large relative correction comes from the region of small $x_L$. The mixing parameter $x_L$ is closely related to the effects of the heavy top quark $T^+$ and $T^-$. When $x_L$ becomes small, the masses of $T^+$ and $T^-$ will become heavy. Due to there are some terms in the amplitudes are proportional to the masses of $T^+$ and $T^-$ even their multi-quadratic terms, which at this time will enhance the relative correction $\delta\sigma/\sigma$.




FIG. 5: The top quark charge asymmetry $A_C(t\bar{t})$ as the function of $f$ for $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV, respectively.

In Fig.5 we show the charge asymmetry $A_C(t\bar{t})$ versus $f$ for the LHC with $\sqrt{s} = 8, 14$ TeV, respectively. In our calculation, the SM prediction of $Ac = 0.0115 \pm 0.0006$[21] has been included. In Fig.5, we give the measurement of the charge asymmetry at the LHC[22] and the $1\sigma$ region. In two cases, we can see the charge asymmetry $A_C(t\bar{t})$ in the LHT model is consistent with the measurement.

In Fig.6 we show the forward-backward asymmetry $A^t_{FB}$ versus $f, x_L$ at the Tevatron. According to the measurements at the Tevatron[23], there is still significant deviation between the measurements and the SM expectation, and disfavor exotic production




FIG. 6: The forward-backward asymmetry in rapidity $A^t_{FB}$ as the function of $f, x_L$ at the Tevatron.

mechanisms with significant negative values[24]. We can see $A^t_{FB}$ in the LHT is negative and very small. So this result can not alleviate the discrepancy between the Tevatron experiment and the SM prediction, and it may give some constraints on the LHT model.

### B. Polarized top quark pair production

(i) The correction to the spin correlation




FIG. 7: The correction to the spin correlation $\delta C$ as the function of $f, x_L$ for $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV, respectively.

In Fig.7 we show the relative correction to the spin correlation $\delta C$ versus $f, x_L$ for the LHC with $\sqrt{s} = 8, 14$ TeV, respectively. Because of the reason similar to

the relative correction of the production cross section $\delta\sigma/\sigma$, we can see $\delta C$ do not decouple obviously with the scale $f$ and the large effects come from the region of small $x_L$. The maximum value of $\delta C$ can reach $-14.5\%$ for $\sqrt{s} = 8$ TeV and $-7.5\%$ for $\sqrt{s} = 14$ TeV, which will be observable at the LHC[25]. Recently, in the helicity basis, the correlation coefficient $\delta C$ is found to be $0.24 \pm 0.02(stat.) \pm 0.08(syst.)$ at the CMS detector of the LHC[26]. The measurements are found to be in agreement with the standard model predictions. So, we can see that the correlation coefficient $\delta C$ will be a good probe for the LHT model at the LHC and may give strong constraints on this model.

(ii) Top quark polarization asymmetry




FIG. 8: The top quark polarization asymmetry $P_t$ as the function of $x_L, M_3$ for $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV, respectively.

In Fig.8 we show the polarization asymmetry $P_t$ versus $f, x_L$ for the LHC with $\sqrt{s} = 8, 14$ TeV, respectively. The maximum value of $P_t$ can respectively reach about $-0.3\%$ for $\sqrt{s} = 8$ TeV and about $-0.6\%$ for $\sqrt{s} = 14$ TeV, which is consistent with the measurement at the LHC[27]. From Fig.8, we can see the large effects come from the region of small $f$ and large $x_L$. Comparing with the results of $\sqrt{s} = 8$ TeV, we can see that the $P_t$ is enhanced greatly for $\sqrt{s} = 14$ TeV.

(iii) Top quark left-right asymmetry

In Fig.9 we show the left-right asymmetry $A_{LR}$ versus $f, x_L$ for the LHC with $\sqrt{s} = 8, 14$ TeV, respectively. We can see that the behavior of $A_{LR}$ for $\sqrt{s} = 14$ TeV




FIG. 9: The top quark left-right asymmetry $A_{LR}$ as the function of $f, x_L$ for $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV, respectively.

is more noticeable than that for $\sqrt{s} = 8$ TeV. The maximum value of $A_{LR}$ can reach about $-1.05\%$ for $\sqrt{s} = 14$ TeV and about $-0.76\%$ for $\sqrt{s} = 8$ TeV, by and large, which is consistent with the SM prediction[28]. In order to analyse the observability of $A_{LR}$ expediently, we follow Refs.[19] and introduce the statistical significance $N_S$. At the LHC with an integrated luminosity $L = 5.0fb^{-1}$, the maximum value of the corresponding $N_S$ for $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV can respectively reach $2.9\sigma$ and $9.3\sigma$. From this result, we can see that the effect of the left-right asymmetry $A_{LR}$ may be detected at the LHC. However, if the effect is not observed in the future, the stronger bounds can be imposed on the LHT model.

### C. The correlation of the observables in $t\bar{t}$ production

In this section, we study the correlations between $\delta C$ and $\delta\sigma/\sigma, A_C(t\bar{t}), P_t, A_{LR}$ at the LHC with $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV, respectively, the corresponding results are displayed in Fig.10. We can see that for the correlation behaviors of the four pairs of quantities there are obviously different between the case $\sqrt{s} = 8$ TeV and the case $\sqrt{s} = 14$ TeV. For the case $\sqrt{s} = 8$ TeV, we can see that the correlations between $\delta C$ and $\delta\sigma/\sigma, A_C(t\bar{t})$ are proportional to each other, respectively. Meanwhile, the correlations between $\delta C$ and $P_t, A_{LR}$ have the similar behavior. For the case $\sqrt{s} = 14$ TeV, we can see that the behaviors of the correlations between $\delta C$ and $\delta\sigma/\sigma, A_C(t\bar{t}), P_t, A_{LR}$ are very

different. According to these correlation behaviors, we can distinguish different models and test the LHT model at the LHC.

## IV. CONCLUSIONS

In the framework of the LHT model, we studied the one-loop effects on the observables in $t\bar{t}$ production and their correlation. We found that the relative correction $\delta\sigma/\sigma$ can reach 4.5% for $\sqrt{s} = 8$ TeV and 2.6% for $\sqrt{s} = 14$ TeV. The relative correction to the spin correlation of top and antitop $\delta C$ can reach $-14.5\%$ for $\sqrt{s} = 8$ TeV and $-7.5\%$ for $\sqrt{s} = 14$ TeV. These effects are significant so that they can be detected at the LHC and can be a probe for the new physics. The charge asymmetry $A_C(t\bar{t})$ and the polarization asymmetries $P_t$, $A_{LR}$ are in agreement with the measurements at the LHC although some of these observables have the opposite sign compared with the SM predictions. The forward-backward asymmetry $A_{\rm FB}^t$ agrees with the SM so that LHT cannot provide an explanation for the observed forward-backward asymmetry at the Tevatron. With the continuous improvement of experimental accuracy, we believed that these results will be tested more strictly.

**Acknowledgments**

We would like to thank Lei Wu for useful discussions and suggestions.

## Appendix: The explicit expressions of the renormalized vertex $\hat{\Gamma}^{\mu}_{gf\bar{f}}$ and the renormalized propagator $-i\hat{\Sigma}^{f}(p)$ [29]

They can be represented in form of 1-point, 2-point and 3-point standard functions $A, B_0, B_1, C_{ij}$. Here $p_t$ and $p_t'$ denote the momenta of the top and antitop respectively, and they are assumed to be outgoing.

(I)Renormalization vertex

$$\hat{\Gamma}^{\mu}_{gf\bar{f}} = \Gamma^{\mu}_{gf\bar{f}} - ieQ_f\gamma^{\mu}(\delta Z_V^f - \gamma_5\delta Z_A^f - \frac{S_W}{2C_W}\delta Z_{ZA}) + ie\gamma^{\mu}(v_f - a_f\gamma_5)\frac{1}{2}\delta Z_{ZA}$$




FIG. 10: The correlations between $\delta C$ and $\delta\sigma/\sigma, A_C(t\bar{t}), P_t, A_{LR}$ for $\sqrt{s} = 8$ TeV and $\sqrt{s} = 14$ TeV, respectively.




where

$$v_f \equiv \frac{I_f^3 - 2Q_f S_W^2}{2C_W S_W}, \quad a_f \equiv \frac{I_f^3}{2C_W S_W}$$

$$\delta Z_{ZA} = 2\frac{\Sigma_T^{AZ}(0)}{M_{Z_L}^2}$$

$$\delta Z_L^f = Re\Sigma_L^f(m_f^2) + m_f^2 \frac{\partial}{\partial P_f^2} Re[\Sigma_L^f(P_f^2) + \Sigma_R^f(P_f^2) + 2\Sigma_S^f(P_f^2)]|_{P_f^2=m_f^2}$$

$$\delta Z_R^f = Re\Sigma_R^f(m_f^2) + m_t^2 \frac{\partial}{\partial P_f^2} Re[\Sigma_L^f(P_f^2) + \Sigma_R^f(P_f^2) + 2\Sigma_S^f(P_f^2)]|_{P_f^2=m_f^2}$$

$$\delta Z_V^f = \frac{1}{2}(\delta Z_L^f + \delta Z_R^f), \delta Z_A^f = \frac{1}{2}(\delta Z_L^f - \delta Z_R^f)$$

$$\hat{\Gamma}^\mu_{u_H\eta} = \frac{g'^2 g_s T^a_{\alpha\beta}(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{100M_{A_H}^2}\frac{i}{16\pi^2}$$
$$\{m_{u_H}^2\gamma^\alpha\gamma^\mu\gamma^\beta C_{\alpha\beta}P_L + m_{u_H}^2(\not{p}_t' + \not{p}_t)\gamma^\mu\gamma^\alpha C_\alpha P_L + m_t^2\gamma^\alpha\gamma^\mu\gamma^\beta C_{\alpha\beta}P_R$$
$$+m_t^2(\not{p}_t' + \not{p}_t)\gamma^\mu\gamma^\alpha C_\alpha P_R - m_{u_H}^2 m_t\gamma^\alpha\gamma^\mu C_\alpha P_R - m_{u_H}^2 m_t(\not{p}_t' + \not{p}_t)\gamma^\mu C_0 P_R$$
$$-m_{u_H}^2 m_t\gamma^\alpha\gamma^\mu C_\alpha P_L - m_{u_H}^2 m_t(\not{p}_t' + \not{p}_t)\gamma^\mu C_0 P_L - m_{u_H}^2 m_t\gamma^\mu\gamma^\alpha C_\alpha P_R$$
$$-m_{u_H}^2 m_t\gamma^\mu\gamma^\alpha C_\alpha P_L + m_{u_H}^4\gamma^\mu C_0 P_L + m_{u_H}^2 m_t^2\gamma^\mu C_0(p_t', p_t, m_{u_H}, m_\eta, m_{u_H})P_R$$
$$+\gamma^\mu[m_t^2 B_1 P_L + m_{u_H}^2 B_1 P_R + \frac{1}{2}(m_t^2 + m_{u_H}^2)B_0(-p_t, m_{u_H}, m_\eta)$$
$$+\frac{1}{2}(m_t^2 + m_{u_H}^2)(m_t^2 + m_{u_H}^2 - m_\eta^2)\frac{\partial}{\partial p_t^2}B_0 - 2m_t^2 m_{u_H}^2\frac{\partial}{\partial p_t^2}B_0]\} \tag{19}$$

$$\hat{\Gamma}^\mu_{u_H\omega^0} = \frac{g^2 g_s T^a_{\alpha\beta}(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{4M_{Z_H}^2}\frac{i}{16\pi^2}$$
$$\{m_{u_H}^2\gamma^\alpha\gamma^\mu\gamma^\beta C_{\alpha\beta}P_L + m_{u_H}^2(\not{p}_t' + \not{p}_t)\gamma^\mu\gamma^\alpha C_\alpha P_L + m_t^2\gamma^\alpha\gamma^\mu\gamma^\beta C_{\alpha\beta}P_R$$
$$+m_t^2(\not{p}_t' + \not{p}_t)\gamma^\mu\gamma^\alpha C_\alpha P_R - m_{u_H}^2 m_t\gamma^\alpha\gamma^\mu C_\alpha P_R - m_{u_H}^2 m_t(\not{p}_t' + \not{p}_t)\gamma^\mu C_0 P_R$$
$$-m_{u_H}^2 m_t\gamma^\alpha\gamma^\mu C_\alpha P_L - m_{u_H}^2 m_t(\not{p}_t' + \not{p}_t)\gamma^\mu C_0 P_L - m_{u_H}^2 m_t\gamma^\mu\gamma^\alpha C_\alpha P_R$$
$$-m_{u_H}^2 m_t\gamma^\mu\gamma^\alpha C_\alpha P_L + m_{u_H}^4\gamma^\mu C_0 P_L + m_{u_H}^2 m_t^2\gamma^\mu C_0(p_t', p_t, m_{u_H}, m_{\omega^0}, m_{u_H})P_R$$
$$+\gamma^\mu[m_t^2 B_1 P_L + m_{u_H}^2 B_1 P_R + \frac{1}{2}(m_t^2 + m_{u_H}^2)B_0(-p_t, m_{u_H}, m_{\omega^0})$$
$$+\frac{1}{2}(m_t^2 + m_{u_H}^2)(m_t^2 + m_{u_H}^2 - m_{\omega^0}^2)\frac{\partial}{\partial p_t^2}B_0 - 2m_t^2 m_{u_H}^2\frac{\partial}{\partial p_t^2}B_0]\} \tag{20}$$

$$\hat{\Gamma}^{\mu}_{d_H\omega^{\pm}} = \frac{g^2 g_s T^a_{\alpha\beta}(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{2M^2_{W_H}} \frac{i}{16\pi^2}$$
$$\{m^2_{d_H}\gamma^{\alpha}\gamma^{\mu}\gamma^{\beta}C_{\alpha\beta}P_L + m^2_{d_H}(\not{p}'_t + \not{p}_t)\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_L + m^2_t\gamma^{\alpha}\gamma^{\mu}\gamma^{\beta}C_{\alpha\beta}P_R$$
$$+m^2_t(\not{p}'_t + \not{p}_t)\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_R - m^2_{d_H}m_t\gamma^{\alpha}\gamma^{\mu}C_{\alpha}P_R - m^2_{d_H}m_t(\not{p}'_t + \not{p}_t)\gamma^{\mu}C_0P_R$$
$$-m^2_{d_H}m_t\gamma^{\alpha}\gamma^{\mu}C_{\alpha}P_L - m^2_{d_H}m_t(\not{p}'_t + \not{p}_t)\gamma^{\mu}C_0P_L - m^2_{d_H}m_t\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_R$$
$$-m^2_{d_H}m_t\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_L + m^4_{d_H}\gamma^{\mu}C_0P_L + m^2_{d_H}m^2_t\gamma^{\mu}C_0P_R(p'_t, p_t, m_{u_H}, m_{\omega^{\pm}}, m_{u_H})$$
$$+\gamma^{\mu}[m^2_tB_1P_L + m^2_{d_H}B_1P_R + \frac{1}{2}(m^2_t + m^2_{d_H})B_0(-p_t, m_{u_H}, m_{\omega^{\pm}})$$
$$+\frac{1}{2}(m^2_t + m^2_{d_H})(m^2_t + m^2_{d_H} - m^2_{\omega^{\pm}})\frac{\partial}{\partial p^2_t}B_0 - 2m^2_tm^2_{d_H}\frac{\partial}{\partial p^2_t}B_0]\} \tag{21}$$

$$\hat{\Gamma}^{\mu}_{T^-\eta} = \frac{4g'^2m^2_t}{25M^2_{A_H}}\frac{f^2}{v^2}[1 - \frac{v^2}{f^2}(\frac{x^2_L}{2} + \frac{1}{6})]^2 g_s T^a_{\alpha\beta}\frac{i}{16\pi^2}$$
$$\{\gamma^{\alpha}\gamma^{\mu}\gamma^{\beta}C_{\alpha\beta}P_R + (\not{p}'_t + \not{p}_t)\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_R + m^2_{T^-}\gamma^{\mu}C_0(p'_t, p_t, m_{T^-}, m_{\eta}, m_{T^-})P_R$$
$$+\gamma_{\mu}[B_1P_L + \frac{1}{2}B_0 + \frac{1}{2}(m^2_t + m^2_{T^-} - m^2_{\eta})\frac{\partial}{\partial p^2_t}B_0(-p_t, m_{T^-}, m_{\eta})]\} \tag{22}$$

$$\hat{\Gamma}^{\mu}_{T^-\omega^0} = \frac{g^2m^2_t}{4M^2_{Z_H}}(\frac{v}{f})^2 g_s T^a_{\alpha\beta}\frac{i}{16\pi^2}$$
$$\{\gamma^{\alpha}\gamma^{\mu}\gamma^{\beta}C_{\alpha\beta}P_R + (\not{p}'_t + \not{p}_t)\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_R + m^2_{T^-}\gamma^{\mu}C_0(p'_t, p_t, m_{T^-}, m_{\omega^0}, m_{T^-})P_R$$
$$+\gamma_{\mu}[B_1P_L + \frac{1}{2}B_0 + \frac{1}{2}(m^2_t + m^2_{T^-} - m^2_{\omega^0})\frac{\partial}{\partial p^2_t}B_0(-p_t, m_{T^-}, m_{\omega^0})]\} \tag{23}$$

$$\hat{\Gamma}^{\mu}_{T^+\pi^0} = \frac{g^2x^2_L}{4M^2_Z\cos^2\theta}(\frac{v}{f})^2 g_s T^a_{\alpha\beta}\frac{i}{16\pi^2}$$
$$\{m^2_{T^+}\gamma^{\alpha}\gamma^{\mu}\gamma^{\beta}C_{\alpha\beta}P_L + m^2_{T^+}(\not{p}'_t + \not{p}_t)\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_L + m^2_t\gamma^{\alpha}\gamma^{\mu}\gamma^{\beta}C_{\alpha\beta}P_R$$
$$+m^2_t(\not{p}'_t + \not{p}_t)\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_R - m^2_{T^+}m_t\gamma^{\alpha}\gamma^{\mu}C_{\alpha}P_R - m^2_{T^+}m_t(\not{p}'_t + \not{p}_t)\gamma^{\mu}C_0P_R$$
$$-m^2_{T^+}m_t\gamma^{\alpha}\gamma^{\mu}C_{\alpha}P_L - m^2_{T^+}m_t(\not{p}'_t + \not{p}_t)\gamma^{\mu}C_0P_L - m^2_{T^+}m_t\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_R$$
$$-m^2_{T^+}m_t\gamma^{\mu}\gamma^{\alpha}C_{\alpha}P_L + m^4_{T^+}\gamma^{\mu}C_0P_L + m^2_{T^+}m^2_t\gamma^{\mu}C_0(p'_t, p_t, m_{T^+}, m_{\pi^0}, m_{T^+})P_R$$
$$+\gamma^{\mu}[m^2_tB_1P_L + m^2_{T^+}B_1P_R + \frac{1}{2}(m^2_t + m^2_{T^+})B_0(-p_t, m_{T^+}, m_{\pi^0})$$
$$+\frac{1}{2}(m^2_t + m^2_{T^+})(m^2_t + m^2_{T^+} - m^2_{\pi^0})\frac{\partial}{\partial p^2_t}B_0 + 2m^2_tm^2_{T^+}\frac{\partial}{\partial p^2_t}B_0]$$
$$+\gamma^{\mu}[m^2_tB_1P_L + m^2_{T^+}B_1P_R + \frac{1}{2}(m^2_t + m^2_{T^+})B_0$$
$$+\frac{1}{2}(m^2_t + m^2_{T^+})(m^2_t + m^2_{T^+} - m^2_{\pi^0})\frac{\partial}{\partial p^2_t}B_0 + 2m^2_tm^2_{T^+}\frac{\partial}{\partial p^2_t}B_0]\} \tag{24}$$

$$\hat{\Gamma}^{\mu}_{T^+h} = m_t^2 g_s T^a_{\alpha\beta}\frac{i}{16\pi^2}$$
$$\{\frac{c_\lambda^2}{s_\lambda^2 v^2}\gamma^\alpha\gamma^\mu\gamma^\beta C_{\alpha\beta}P_L + \frac{c_\lambda^2}{s_\lambda^2 v^2}(\not{p}'_t+\not{p}_t)\gamma^\mu\gamma^\alpha C_\alpha P_L + \frac{s_\lambda^4}{f^2}\gamma^\alpha\gamma^\mu\gamma^\beta C_{\alpha\beta}P_R$$
$$+\frac{s_\lambda^4}{f^2}(\not{p}'_t+\not{p}_t)\gamma^\mu\gamma^\alpha C_\alpha P_R - \frac{c_\lambda m_{T^+}s_\lambda}{vf}\gamma^\alpha\gamma^\mu C_\alpha P_R - \frac{c_\lambda m_{T^+}s_\lambda}{vf}(\not{p}'_t+\not{p}_t)\gamma^\mu C_0 P_R$$
$$-\frac{c_\lambda m_{T^+}s_\lambda}{vf}\gamma^\alpha\gamma^\mu C_\alpha P_L - \frac{c_\lambda m_{T^+}s_\lambda}{vf}(\not{p}'_t+\not{p}_t)\gamma^\mu C_0 P_L - \frac{c_\lambda m_{T^+}s_\lambda}{vf}\gamma^\mu\gamma^\alpha C_\alpha P_R$$
$$-\frac{c_\lambda m_{T^+}s_\lambda}{vf}\gamma^\mu\gamma^\alpha C_\alpha P_L + \frac{c_\lambda^2 m_{T^+}^2}{s_\lambda^2 v^2}\gamma^\mu C_0 P_L + m_{T^+}^2\frac{s_\lambda^4}{f^2}\gamma^\mu C_0(p'_t,p_t,m_{T^+},m_h,m_{T^+})P_R$$
$$+\gamma^\mu[\frac{s_\lambda^4}{f^2}B_1 P_L + \frac{c_\lambda^2}{s_\lambda^2 v^2}B_1 P_R + \frac{1}{2}(\frac{s_\lambda^4}{f^2}+\frac{c_\lambda^2}{s_\lambda^2 v^2})B_0(-p_t,m_{T^+},m_h)$$
$$+\frac{1}{2}(\frac{s_\lambda^4}{f^2}+\frac{c_\lambda^2}{s_\lambda^2 v^2})(m_t^2+m_{T^+}^2-m_h^2)\frac{\partial}{\partial p_t^2}B_0 - 2\frac{s_\lambda^4}{f^2}m_{T^+}^2\frac{\partial}{\partial p_t^2}B_0]\} \quad (25)$$

$$\hat{\Gamma}^{\mu}_{u_H A_H} = \frac{g'^2 g_s T^a_{\alpha\beta}(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{100}\frac{i}{16\pi^2}\{2\gamma^\sigma\gamma^\mu\gamma^\lambda C_{\lambda\sigma}P_L$$
$$+2\gamma^\mu P_L - \gamma^\rho(\not{p}'_t+\not{p}_t)\gamma^\mu\gamma^\lambda\gamma_\rho C_\lambda P_L + 2m_{u_H}^2\gamma^\mu C_0(p'_t,p_t,m_{u_H},M_{A_H},m_{u_H})P_L$$
$$+\gamma^\mu[2B_1 P_R + B_0 + (m_t^2+m_{u_H}^2-M_{A_H}^2)\frac{\partial}{\partial p_t^2}B_0(-p_t,m_{u_H},M_{A_H}) - P_L]\} \quad (26)$$

$$\hat{\Gamma}^{\mu}_{u_H Z_H} = \frac{g^2 g_s T^a_{\alpha\beta}(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{4}\frac{i}{16\pi^2}\{2\gamma^\sigma\gamma^\mu\gamma^\lambda C_{\lambda\sigma}P_L$$
$$+2\gamma^\mu P_L - \gamma^\rho(\not{p}'_t+\not{p}_t)\gamma^\mu\gamma^\lambda\gamma_\rho C_\lambda P_L + 2m_{u_H}^2\gamma^\mu C_0(p'_t,p_t,m_{u_H},M_{Z_H},m_{u_H})P_L$$
$$+\gamma^\mu[2B_1 P_R + B_0 + (m_t^2+m_{u_H}^2-M_{Z_H}^2)\frac{\partial}{\partial p_t^2}B_0(-p_t,m_{u_H},M_{Z_H}) - P_L]\} \quad (27)$$

$$\hat{\Gamma}^{\mu}_{d_H W_H} = \frac{g^2 g_s T^a_{\alpha\beta}(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{2}\frac{i}{16\pi^2}\{2\gamma^\sigma\gamma^\mu\gamma^\lambda C_{\lambda\sigma}P_L$$
$$+2\gamma^\mu P_L - \gamma^\rho(\not{p}'_t+\not{p}_t)\gamma^\mu\gamma^\lambda\gamma_\rho C_\lambda P_L + 2m_{d_H}^2\gamma^\mu C_0(p'_t,p_t,m_{d_H},M_{W_H},m_{d_H})P_L$$
$$+\gamma^\mu[2B_1 P_R + B_0 + (m_t^2+m_{d_H}^2-M_{W_H}^2)\frac{\partial}{\partial p_t^2}B_0(-p_t,m_{d_H},M_{W_H}) - P_L]\} \quad (28)$$

$$\hat{\Gamma}^{\mu}_{T^- A_H} = \frac{4g'^2 g_s T^a_{\alpha\beta}}{25}\frac{i}{16\pi^2}$$
$$\{2\gamma^\sigma\gamma^\mu\gamma^\lambda x_L C_{\lambda\sigma}P_R + 2x_L\gamma^\mu P_R + 2\gamma^\lambda\gamma^\mu(\not{p}'_t+\not{p}_t)x_L C_\lambda(p'_t,p_t,m_{T^-},M_{A_H},m_{T^-})P_R$$
$$-8m_{T^-}x_L\sqrt{x_L}\frac{v}{f}C^\mu - 4(p'_t+p_t)^\mu m_{T^-}x_L\sqrt{x_L}\frac{v}{f}C_0 + 2m_{T^-}^2 x_L\gamma^\mu C_0 P_R$$
$$+\gamma^\mu[2x_L B_1 P_L + x_L B_0 + x_L(m_t^2+m_{T^-}^2-M_{A_H}^2)\frac{\partial}{\partial p_t^2}B_0 - P_L$$
$$-8x_L\sqrt{x_L}\frac{v}{f}m_t m_{T^-}\frac{\partial}{\partial p_t^2}B_0 - \frac{\partial}{\partial p_t^2}B_0(-p_t,m_{T^-},M_{A_H}) - x_L P_R]\} \quad (29)$$

$$\hat{\Gamma}^{\mu}_{T^+Z} = \frac{g^2 g_s T^a_{\alpha\beta} x_L^2}{4\cos^2\theta}\frac{v^2}{f^2}\frac{i}{16\pi^2}\{2\gamma^\sigma\gamma^\mu\gamma^\lambda C_{\lambda\sigma}P_L + 2\gamma^\mu P_L$$
$$-\gamma^\rho(\not{p}'_t + \not{p}_t)\gamma^\mu\gamma^\lambda\gamma_\rho C_\lambda P_L + 2m^2_{T^+}\gamma^\mu C_0(p'_t, p_t, m_{T^+}, M_Z, m_{T^+})P_L$$
$$+\gamma^\mu[4B_1 P_R + 2B_0 + 2(m_t^2 + m^2_{T^+} - m_Z^2)\frac{\partial}{\partial p_t^2}B_0(-p_t, m_{T^+}, M_Z) - 2P_L]\} \qquad (30)$$

(II)Renormalization fermion propagator




$$-i\hat{\Sigma}^f(p) = -i\Sigma^f(p) + (-i\delta\Sigma^f(p))$$

where

$$\Sigma^f(p) = m_f\Sigma_S^f(p^2) + \not{p}P_L\Sigma_L^f(p^2) + \not{p}P_R\Sigma_R^f(p^2)$$
$$\delta\Sigma^f(p) = \delta m_f + m_f\frac{1}{2}\delta Z_L^f + m_f\frac{1}{2}\delta Z_R^f - \not{p}P_L\delta Z_L^f - \not{p}P_R\delta Z_R^f$$
$$\delta m_f = -m_f Re[\Sigma_S^f(m_f^2) + \frac{1}{2}\Sigma_L^f(m_f^2) + \frac{1}{2}\Sigma_R^f(m_f^2)]$$
$$\delta Z_L^f = Re\Sigma_L^f(m_f^2) + m_f^2\frac{\partial}{\partial p^2}Re[\Sigma_L^f(p^2) + \Sigma_R^f(p^2) + 2\Sigma_S^f(p^2)]|_{p^2=m_f^2}$$
$$\delta Z_R^f = Re\Sigma_R^f(m_f^2) + m_f^2\frac{\partial}{\partial p^2}Re[\Sigma_L^f(p^2) + \Sigma_R^f(p^2) + 2\Sigma_S^f(p^2)]|_{p^2=m_f^2}$$

$$-i\hat{\Sigma}^{\eta u_H}(p) = -\frac{g'^2(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{100M^2_{A_H}}\frac{i}{16\pi^2}$$
$$\{m^2_{u_H}\not{p}_t B_1 P_L + m_t^2\not{p}_t B_1 P_R + m_t m^2_{u_H}B_0(-p_t, m_{u_H}, m_\eta)$$
$$-m_t m^2_{u_H}B_0 + m_t^3\frac{\partial}{\partial p_t^2}(m^2_{u_H}B_1 + m_t^2 B_1 + 2m^2_{u_H}B_0)$$
$$-\not{p}_t P_L[m^2_{u_H}B_1 + m_t^2\frac{\partial}{\partial p_t^2}(m^2_{u_H}B_1 + m_t^2 B_1 + 2m^2_{u_H}B_0)]$$
$$-\not{p}_t P_R[m^2_{u_H}B_1 + m_t^2\frac{\partial}{\partial p_t^2}(m^2_{u_H}B_1 + m_t^2 B_1 + 2m^2_{u_H}B_0)]\} \qquad (31)$$

$$-i\hat{\Sigma}^{\omega^0 u_H}(p) = -\frac{g'^2(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{100M^2_{A_H}}\frac{i}{16\pi^2}$$
$$\{m^2_{u_H}\not{p}_t B_1 P_L + m_t^2\not{p}_t B_1 P_R + m_t m^2_{u_H}B_0(-p_t, m_{u_H}, m_{\omega^0})$$
$$-m_t m^2_{u_H}B_0 + m_t^3\frac{\partial}{\partial p_t^2}(m^2_{u_H}B_1 + m_t^2 B_1 + 2m^2_{u_H}B_0)$$
$$-\not{p}_t P_L[m^2_{u_H}B_1 + m_t^2\frac{\partial}{\partial p_t^2}(m^2_{u_H}B_1 + m_t^2 B_1 + 2m^2_{u_H}B_0)]$$
$$-\not{p}_t P_R[m^2_{u_H}B_1 + m_t^2\frac{\partial}{\partial p_t^2}(m^2_{u_H}B_1 + m_t^2 B_1 + 2m^2_{u_H}B_0)]\} \qquad (32)$$

$$-i\hat{\Sigma}^{\omega^{\pm}d_H}(p) = -\frac{g'^2(V_{Hu})^*_{i3}(V_{Hu})_{i3}}{100M^2_{A_H}}\frac{i}{16\pi^2}$$
$$\{m^2_{d_H}\not{p}_t B_1 P_L + m^2_t \not{p}_t B_1 P_R + m_t m^2_{d_H} B_0(-p_t, m_{d_H}, m_{\omega^{\pm}})$$
$$-m_t m^2_{d_H} B_0 + m^3_t \frac{\partial}{\partial p^2_t}(m^2_{d_H} B_1 + m^2_t B_1 + 2m^2_{d_H} B_0)$$
$$-\not{p}_t P_L[m^2_{d_H} B_1 + m^2_t \frac{\partial}{\partial p^2_t}(m^2_{d_H} B_1 + m^2_t B_1 + 2m^2_{d_H} B_0)]$$
$$-\not{p}_t P_R[m^2_{d_H} B_1 + m^2_t \frac{\partial}{\partial p^2_t}(m^2_{d_H} B_1 + m^2_t B_1 + 2m^2_{d_H} B_0)]\} \tag{33}$$

$$-i\hat{\Sigma}^{\eta T^-}(p) = -(\frac{2g' m_t f}{5M_{A_H} v})^2 \frac{i}{16\pi^2}[1 - \frac{v^2}{f^2}(\frac{x^2_L}{2} + \frac{1}{6})]^2[\not{p}_t B_1 P_R$$
$$+m^3_t \frac{\partial}{\partial p^2_t} B_1 - \not{p}_t P_L m^2_t \frac{\partial}{\partial p^2_t} B_1 - \not{p}_t P_R (B_1 + m^2_t \frac{\partial}{\partial p^2_t} B_1)(-p_t, m_{T^-}, m_\eta)] \tag{34}$$

$$-i\hat{\Sigma}^{\omega_0 T^-}(p) = -(\frac{g m_t f}{2M_{Z_H} v})^2 \frac{i}{16\pi^2}$$
$$[\not{p}_t B_1 P_R + m^3_t \frac{\partial}{\partial p^2_t} B_1 - \not{p}_t P_L m^2_t \frac{\partial}{\partial p^2_t} B_1 - \not{p}_t P_R B_1(-p_t, m_{T^-}, m_{\omega_0})] \tag{35}$$

$$-i\hat{\Sigma}^{\pi^0 T^+}(p) = -\frac{g^2 x^2_L v^2}{4M^2_Z \cos^2\theta f^2}\frac{i}{16\pi^2}$$
$$\{m^2_{T^+}\not{p}_t B_1 P_L + m^2_t \not{p}_t B_1 P_R + m_t m^2_{T^+} B_0(-p_t, m_{T^+}, m_{\pi^0})$$
$$-m_t m^2_{T^+} B_0 + m^3_t \frac{\partial}{\partial p^2_t}(m^2_{T^+} B_1 + m^2_t B_1 + 2m^2_{T^+} B_0)$$
$$-\not{p}_t P_L[m^2_{T^+} B_1 + m^2_t \frac{\partial}{\partial p^2_t}(m^2_{T^+} B_1 + m^2_t B_1 + 2m^2_{T^+} B_0)]$$
$$-\not{p}_t P_R[m^2_{T^+} B_1 + m^2_t \frac{\partial}{\partial p^2_t}(m^2_{T^+} B_1 + m^2_t B_1 + 2m^2_{T^+} B_0)]\} \tag{36}$$

$$-i\hat{\Sigma}^{hT^+}(p) = m^2_t \frac{i}{16\pi^2}\{\frac{x_L}{(1-x_L)v^2}\not{p}_t B_1 P_L - \frac{(1-x_L)^2}{f^2}\not{p}_t B_1 P_R$$
$$-\frac{s_\lambda\sqrt{x_L} m_{T^+}}{fv} B_0 + \frac{s_\lambda\sqrt{x_L} m_{T^+}}{fv} B_0(-p_t, m_{T^+}, m_h)$$
$$+m_t \frac{\partial}{\partial p^2_t}[\frac{x_L m^2_t}{(1-x_L)v^2} B_1 - \frac{(1-x_L)^2 m^2_t}{f^2} B_1 - \frac{2s_\lambda\sqrt{x_L} m_{T^+} m_t}{fv} B_0] \tag{37}$$
$$-\not{p}_t P_L[\frac{x_L}{(1-x_L)v^2} B_1 + \frac{\partial}{\partial p^2_t}[\frac{x_L m^2_t}{(1-x_L)v^2} B_1 - \frac{(1-x_L)^2 m^2_t}{f^2} B_1 - \frac{2s_\lambda\sqrt{x_L} m_{T^+} m_t}{fv} B_0]$$
$$-\not{p}_t P_R[\frac{x_L}{(1-x_L)v^2} B_1 + \frac{\partial}{\partial p^2_t}[\frac{x_L m^2_t}{(1-x_L)v^2} B_1 - \frac{(1-x_L)^2 m^2_t}{f^2} B_1 - \frac{2s_\lambda\sqrt{x_L} m_{T^+} m_t}{fv} B_0]\}$$

$$-i\hat{\Sigma}^{A_H u_H}(p) = -\frac{g'^2 (V_{Hu})^*_{i3}(V_{Hu})_{i3}}{100}\frac{i}{16\pi^2}[2\not{p}_t(B_1+\frac{1}{2})P_L$$
$$+2m_t^3\frac{\partial}{\partial p_t^2}B_1 - 2\not{p}_t P_L(B_1+\frac{1}{2}+m_t^2\frac{\partial}{\partial p_t^2}B_1) - 2\not{p}_t P_R m_t^2\frac{\partial}{\partial p_t^2}B_1(-p_t, m_{u_H}, M_{A_H})] \quad (38)$$

$$-i\hat{\Sigma}^{Z_H u_H}(p) = -\frac{g^2 (V_{Hu})^*_{i3}(V_{Hu})_{i3}}{4}\frac{i}{16\pi^2}[2\not{p}_t(B_1+\frac{1}{2})P_L$$
$$+2m_t^3\frac{\partial}{\partial p_t^2}B_1 - 2\not{p}_t P_L(B_1+\frac{1}{2}+m_t^2\frac{\partial}{\partial p_t^2}B_1) - 2\not{p}_t P_R m_t^2\frac{\partial}{\partial p_t^2}B_1(-p_t, m_{u_H}, M_{Z_H})] \quad (39)$$

$$-i\hat{\Sigma}^{W_H d_H}(p) = -\frac{g^2 (V_{Hu})^*_{i3}(V_{Hu})_{i3}}{2}\frac{i}{16\pi^2}[2\not{p}_t(B_1+\frac{1}{2})P_L$$
$$+2m_t^3\frac{\partial}{\partial p_t^2}B_1 - 2\not{p}_t P_L(B_1+\frac{1}{2}+m_t^2\frac{\partial}{\partial p_t^2}B_1) - 2\not{p}_t P_R m_t^2\frac{\partial}{\partial p_t^2}B_1(-p_t, m_{d_H}, M_{W_H})] \quad (40)$$

$$-i\hat{\Sigma}^{ZT^+}(p) = -\frac{g^2 x_L^2 v^2}{4\cos^2\theta f^2}\frac{i}{16\pi^2}[2\not{p}_t(B_1+\frac{1}{2})P_L$$
$$+2m_t^3\frac{\partial}{\partial p_t^2}B_1 - 2\not{p}_t P_L(B_1+\frac{1}{2}+m_t^2\frac{\partial}{\partial p_t^2}B_1) - 2\not{p}_t P_R m_t^2\frac{\partial}{\partial p_t^2}B_1(-p_t, m_{T^+}, M_Z)] \quad (41)$$

$$-i\hat{\Sigma}^{A_H T^-}(p) = -\frac{4g'^2}{25}\frac{i}{16\pi^2}\{2x_L^2\frac{v^2}{f^2}\not{p}_t(B_1+\frac{1}{2})P_L + 2x_L\not{p}_t(B_1+\frac{1}{2})P_R$$
$$+2m_t^3\frac{\partial}{\partial p_t^2}[x_L^2\frac{v^2}{f^2}B_1 + x_L B_1 + 4x_L\sqrt{x_L}\frac{v}{f}\frac{m_{T^-}}{m_t}B_0(-p_t, m_{T^-}, M_{A_H})]$$
$$-2\not{p}_t P_L[x_L^2\frac{v^2}{f^2}(B_1+\frac{1}{2}) + m_t^2\frac{\partial}{\partial p_t^2}(x_L^2\frac{v^2}{f^2}B_1 + x_L B_1 + 4x_L\sqrt{x_L}\frac{v}{f}\frac{m_{T^-}}{m_t}B_0)]$$
$$-2\not{p}_t P_R[x_L^2\frac{v^2}{f^2}(B_1+\frac{1}{2}) + m_t^2\frac{\partial}{\partial p_t^2}(x_L^2\frac{v^2}{f^2}B_1 + x_L B_1 + 4x_L\sqrt{x_L}\frac{v}{f}\frac{m_{T^-}}{m_t}B_0)]\} \quad (42)$$

(III)Renormalization gluon propagator




$$-i\hat{\Sigma}_{\mu\nu}(k) = -i\Sigma_{\mu\nu}(k) + (-i\delta\Sigma_{\mu\nu}(k))$$

where

$$\Sigma_{\mu\nu}(k) = g_{\mu\nu}\Sigma_T(k) + k_\mu k_\nu \Sigma_L(k)$$
$$\delta\Sigma^{gg}_{\mu\nu}(k) = g_{\mu\nu}[\delta Z_{gg} k^2]$$
$$\delta Z_{gg} = -\frac{\partial\Sigma_T^{gg}(k^2)}{\partial k^2}|_{k^2=0}$$
$$\hat{\Sigma}^{gg}_{\mu\nu} = \hat{\Sigma}^{gg}_{\mu\nu}(u_H) + \hat{\Sigma}^{gg}_{\mu\nu}(d_H) + \hat{\Sigma}^{gg}_{\mu\nu}(T^\pm)$$

$$
\begin{aligned}
&-i\hat{\Sigma}^{gg}_{\mu\nu}(k) = \frac{-4ig_s^2 g_{\mu\nu} T^a_{\alpha\beta} T^b_{\beta\alpha}}{16\pi^2} \\
&\{[-\frac{2}{3}A_0 + \frac{2}{3}m^2_{T^-}B_0 - \frac{2}{3}k^2 B_1(k, m_{T^-}, m_{T^-}) + \frac{2}{3}m^2_{T^-} - \frac{k^2}{9}] \\
&+[-\frac{2}{3}A_0 + \frac{2}{3}m^2_{T^+}B_0 - \frac{2}{3}k^2 B_1(k, m_{T^+}, m_{T^+}) + \frac{2}{3}m^2_{T^+} - \frac{k^2}{9}] \\
&+[-\frac{2}{3}A_0 + \frac{2}{3}m^2_{u_H}B_0 - \frac{2}{3}k^2 B_1(k, m_{u_H}, m_{u_H}) + \frac{2}{3}m^2_{u_H} - \frac{k^2}{9}] \\
&+[-\frac{2}{3}A_0 + \frac{2}{3}m^2_{d_H}B_0 - \frac{2}{3}k^2 B_1(k, m_{d_H}, m_{d_H}) + \frac{2}{3}m^2_{d_H} - \frac{k^2}{9}] \\
&-[\frac{2}{3}m^2_{T^-}\frac{\partial B_0}{\partial k^2} - \frac{2}{3}B_1(0, m_{T^-}, m_{T^-}) - \frac{1}{9}]k^2 \\
&-[\frac{2}{3}m^2_{T^+}\frac{\partial B_0}{\partial k^2} - \frac{2}{3}B_1(0, m_{T^+}, m_{T^+}) - \frac{1}{9}]k^2 \\
&-[\frac{2}{3}m^2_{u_H}\frac{\partial B_0}{\partial k^2} - \frac{2}{3}B_1(0, m_{u_H}, m_{u_H}) - \frac{1}{9}]k^2 \\
&-[\frac{2}{3}m^2_{d_H}\frac{\partial B_0}{\partial k^2} - \frac{2}{3}B_1(0, m_{d_H}, m_{d_H}) - \frac{1}{9}]k^2\}
\end{aligned} \tag{43}
$$

---


[1] G. Stricker et al., CDF note 9724 (2009); T. Aaltonen et al. [CDF Collaboration], Phys. Rev. Lett. 101, 202001 (2008); V. M. Abazov et al. [D0 Collaboration], Phys. Rev. Lett. 100, 142002 (2008).

[2] A. Djouadi,G. Moreau,F. Richard, R.K. Singh, Phys.Rev.D 82:071702 (2010); S. Jung, H. Murayama, A. Pierce,J.D. Wells,Phys.Rev.D 81:015004 (2010); J. Shu,T.M.P. Tait, K. Wang, Phys.Rev.D 81:034012 (2010); A. Arhrib, R. Benbrik and C. H. Chen, Phys.Rev. D82 034034(2010); D. W. Jung, P. Ko, J. S. Lee and S. h. Nam, Phys. Lett. B 691, 238 (2010); I. Dorsner, S. Fajfer, J.F. Kamenik,N. Kosnik, Phys.Rev.D 81:055009(2010); V. Barger, W.-Y. Keung, C.-T. Yu, Phys. Rev. D 81, 113009 (2010); C. H. Chen, G. Cvetic and C. S. Kim, Phys.Lett.B694:393-397(2011); M. V. Martynov and A. D. Smirnov, Mod. Phys. Lett. A 25, 2637 (2010); J. A. Aguilar-Saavedra, Nucl. Phys. B 843, 638 (2011); C. Degrande, et al., JHEP 1103:125(2011); B. Xiao,Y.K. Wang,S.H. Zhu, Phys.Rev.D 82:034026(2010); S. Jung, H. Murayama, A. Pierce and J. D. Wells, Phys. Rev. D 81, 015004 (2010);J. Cao,Z. Heng,L. Wu,J.M. Yang, Phys. Rev. D 81, 014016 (2010); P. H. Frampton, J. Shu and K. Wang, Phys.Lett.B683:294-297(2010); R. S. Chivukula, E. H. Simmons, C.-P. Yuan, Phys.Rev.D82:094009(2010); Junjie Cao, Lin Wang, Lei Wu, Jin Min Yang Phys.Rev. D

84: 074001(2011).

[3] W. Bernreuther, J. Phys. G35, 083001,(2008) D. Chakraborty, J. Konigsberg, D. Rainwater, Ann. Rev. Nucl. Part. Sci. 53, 301 (2003); E. H. Simmons, hep-ph/0211335; C.-P. Yuan, hep-ph/0203088; S. Willenbrock, hep-ph/0211067; M. Beneke, et al., hep-ph/0003033; T. Han, arXiv:0804.3178; C. T. Hill and S. J. Parke, Phys. Rev. D 49, 4454 (1994); K. Whisnant,J.M. Yang, B.L.Young,X. Zhang, Phys. Rev. D 56, 467 (1997); J. M. Yang, B.-L. Young, Phys. Rev. D 56, 5907 (1997); K.I. Hikasa, K. Whisnant,J.M. Yang,B.L. Young, Phys. Rev. D 58, 114003 (1998); R.A. Coimbra, et al., Phys.Rev.D79:014006,2009; Jin-Yan Liu, Zong-Guo Si, Chong-Xing Yue, Phys.Rev. D81 (2010) 015011.

[4] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, Phys. Lett. B 513, 232 (2001); N. Arkani-Hamed, et al., JHEP 0208, 020 (2002); JHEP 0208, 021 (2002); I. Low, W. Skiba, and D.Smith, Phys. Rev. D 66, 072001 (2002); D. E. Kaplan and M. Schmaltz, JHEP 0310, 039(2003).

[5] N. Arkani-Hamed, A. G. Cohen, E. Katz, and A. E. Nelson, JHEP 0207, 034 (2002); S.Chang, JHEP 0312, 057 (2003); T. Han, H. E. Logan, B. McElrath, and L. T. Wang, Phys.Rev. D 67, 095004 (2003); M. Schmaltz and D. Tucker-smith, Ann. Rev. Nucl. Part. Sci. 55,229 (2005).

[6] C.Csaki, J.Hubisz, G.D.Kribs, P.Meade, J.Terning, Phys. Rev. D 67, 115002 (2003); Phys. Rev. D 68, 035009 (2003); J. L. Hewett,F. J. Petriello, and T. G. Rizzo, JHEP 0310, 062 (2003); M. C. Chen and S. Dawson, Phys.Rev. D 70, 015003 (2004); M. C. Chen, et al., Mod. Phys. Lett. A 21, 621 (2006); W. Kilian, J. Reuter, Phys. Rev. D 70, 015004 (2004).

[7] G. Marandella, C. Schappacher and A. Strumia, Phys. Rev. D 72, 035014 (2005).

[8] H. C. Cheng and I. Low, JHEP 0309, 051 (2003); JHEP 0408, 061 (2004); I. Low, JHEP0410, 067 (2004); J. Hubisz and P. Meade, Phys. Rev. D 71, 035016 (2005).

[9] J. H. Kuhn, Nucl. Phys. B 237, 77 (1984); V. D. Barger, J. Ohnemus and R. J. N. Phillips, Int. J. Mod. Phys. A 4, 617 (1989); G. Mahlon and S. J. Parke, Phys. Lett. B 411, 173 (1997); Phys. Rev. D 53, 4886 (1996); Phys. Rev. D 81, 074024 (2010); T. Stelzer and S. Willenbrock, Phys. Lett. B 374, 169 (1996); Bernreuther,W.,Brandenburg,A.,Si,Z.G.,Uwer,P., Phys. Rev. Lett. 87, 242002 (2001); W. Bernreuther and Z. G. Si, Nucl. Phys. B 837, 90 (2010); G. Mahlon, arXiv:1007.1716 [hep-ph].

[10] K. Cheung, Phys. Rev. D 55, 4430 (1997); B. Holdom and T. Torma, Phys. Rev. D 60, 114010 (1999); M. Arai,N. Okada,K. Smolek,V. Simak, Phys. Rev. D 70, 115015 (2004); M. Arai,N. Okada,K. Smolek,V. Simak, Phys. Rev. D 75, 095008 (2007); Acta Phys. Polon. B 40, 93 (2009); M. Arai, N. Okada and K. Smolek, Phys. Rev. D 79, 074019 (2009); C. X. Yue, T. T. Zhang and J. Y. Liu, J. Phys. G 37, 075016 (2010); Chengcheng Han, Ning Liu, Lei Wu, Jin Min Yang, Phys.Lett. B714 (2012) 295-300; Lei Wang, Lei Wu, Jin Min Yang, Phys.Rev. D85 (2012) 075017; Junjie Cao, Lei Wu, Jin Min Yang,Phys.Rev. D83 (2011) 034024.

[11] Jay Hubisz, Seung J. Lee, Gil Paz, JHEP0606:041 (2006); M.Blanke, et al., Phys. Lett. B 646, 253 (2007).

[12] M.Blanke, et al., Phys. Lett. B 646, 253 (2007); M.Blanke, et al., JHEP 0701, 066 (2007).

[13] H.L. Lai, M. Guzzi, J. Huston,Z. Li,P.M. Nadolsky, J. Pumplin,C.P.Yuan, Phys.Rev.D 82, 074024 (2010).

[14] K. Nakamura et al. (Particle Data Group), J. Phys. G 37 (2010) 075021.

[15] G. Aad et al. [ATLAS Collaboration], Phys.Lett. B716 (2012) 1-29, arXiv:1207.7214; S. Chatrchyan et al. [CMS Collaboration], CMS-HIG-12-028, arXiv:1207.7235.

[16] M.Blanke, et al.,JHEP 0705:013(2007).

[17] J. Hubisz, P. Meade, A. Noble, and M. Perelstein, J. High Energy Phys. 0601 (2006) 135; A. Belyaev, C.-R. Chen, K. Tobe, and C.P. Yuan, Phys. Rev. D 74, 115020 (2006); Q.-H. Cao and C.-R. Chen,Phys. Rev. D 76, 075007 (2007); Luca Fiorini, ATL-PHYS-PROC-2012-029, arXiv:1201.5844.

[18] J.H. Kuhn,G. Rodrigo,Phys. Rev. D 59, 054017 (1999); M. T. Bowen, S. D. Ellis and D. Rainwater, Phys. Rev. D 73, 014008 (2006); V. Ahrens et al., JHEP 1009, 097 (2010); N. Kidonakis, Phys.Rev.D84:011504,2011; W. Hollik and D. Pagani, Phys.Rev. D84 (2011) 093003.

[19] C. Kao and D. Wackeroth, Phys. Rev. D 61, 055009 (2000); K.I. Hikasa,J.M. Yang,B.L. Young,Phys. Rev. D 60, 114041 (1999); P. Y. Li, et al., Eur. Phys. J. C 51, 163 (2007); David Krohn, Tao Liu, Jessie Shelton, Lian-Tao Wang, Phys.Rev. D84 (2011) 074034.

[20] Qing-Hong Cao, Chuan-Ren Chen, F. Larios, C.-P. Yuan, Phys.Rev.D79:015004,2009.

[21] Johann H. Kuhn, German Rodrigo,arXiv:1109.6830.

[22] CMS Collaboration, Phys. Lett. B 709 (2012) 28-49; The ATLAS Collaboration, Eur.Phys.J. C72 (2012) 2039.

[23] http://www-cdf.fnal.gov/physics/new/top/2011/AfbComb/.

[24] T. Aaltonen, et al, CDF Collaboration, Phys.Rev.Lett.101:202001,2008.

[25] F. Hubaut, et al., Eur. Phys. J. C 44 13 (2005).

[26] https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsTOP12004; https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/CONFNOTES/ATLAS-CONF-2011-117/.

[27] https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsTOP12016.

[28] C. Kao and D. Wackeroth, Phys. Rev. D 61, 055009 (2000).

[29] W.F.L.Hollik, Fortschr. Phys. 38: 165-260(1990); A.Denner, Fortschr. Phys.41: 307-420 (1993).